


*Considerations on the suppression
of charged particles
in high energy heavy ion collisions*

Mihai Petrovici, Amelia Lindner and Amalia Pop

HADRON PHYSICS DEPARTMENT









Outline

- **Introduction**
- **R_{AA} , $\langle dN/dy \rangle / S_{\perp}$ - $\langle N_{\text{part}} \rangle$, $\langle dN_{\text{ch}}/d\eta \rangle$ dependence**
 - **Cu-Cu, Au-Au - $\sqrt{s_{NN}} = 200$ GeV;**
 - **Pb-Pb - $\sqrt{s_{NN}} = 2.76$ and 5.02 TeV;**
 - **Xe-Xe - $\sqrt{s_{NN}} = 5.44$ TeV**
 - **Core-Corona contribution**
 - **Suppression saturation at LHC energies**
- **R_{AA}^N ($\langle N_{\text{bin}} \rangle \rightarrow \langle dN_{\text{ch}}/d\eta \rangle^{A-A, \text{cen}} / \langle dN_{\text{ch}}/d\eta \rangle^{\text{pp, MB}}$) - $\langle N_{\text{part}} \rangle$, $\langle dN_{\text{ch}}/d\eta \rangle$ dependence**
 - **Considerations on the missing suppression in high charged particle multiplicity events for pp collisions at $\sqrt{s} = 7$ TeV**
- **R_{CP} , R_{CP}^N - $\langle N_{\text{part}} \rangle$ dependence**
- **charged particles - R_{CP} , R_{CP}^N and π^0 - R_{AA} , R_{AA}^N as a function of $\sqrt{s_{NN}}$**
- **Outlook**

Suppression as a tomographic probe for deconfined matter studies


- Proposed by Bjorken ~ 40 years ago for hadron-hadron collisions
FERMILAB-PUB-82-059-THY, 1982
- confirmed in heavy ion collisions by many experiments, for different energies and system size
- interpreted and estimated by many theoretical models

(not yet confirmed)



$$Ratio = \frac{[\frac{d^2N}{dydp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]_{cen,mult}}{[\frac{d^2N}{dydp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]_{pp,MB}}$$

J. Liao and E. Shuryak, PRL 102(2009)202302





JET Collaboration, Phys. Rev. C90(2014)014909 and references therein

J.Liao, S.Shi and M.Gyulassy. Chin. Phys. C43(209)044101

However, a proper description of the parton energy loss in the non-equilibrium expanding deconfined matter for the intermediate p_T range remains a challenging task

Centrality, $\langle N_{part} \rangle$, $\langle dN_{ch}/d\eta \rangle$ correlations

(based on the Glauber MC approach)



- PHOBOS Collaboration, *PRL* 96(2006)212301
- STAR Collaboration, *PRL* 91(2003)172302
- ALICE Collaboration, *Phys.Lett. B* 788(2019)166
- ALICE Collaboration, *PRL* 116(2016)222302
- M.Petrovici et al. *Phys. Rev. C* 98(2018)024904



- STAR Collaboration, *Phys. Rev. C* 79(2009)034909
- PHENIX Collaboration, *Phys. Rev. C* 93(2016)024901
- ALICE Collaboration, *Phys. Rev. C* 88(2013)044910

Transverse overlapping area

Percentage of nucleons undergoing a single collision (Corona)

(based on the Glauber MC approach)

$$S_{\perp}^{\text{var}} \sim (\langle \sigma_x^2 \rangle \langle \sigma_y^2 \rangle - \langle \sigma_{xy} \rangle^2)^{1/2}$$



- $S_{\perp}^{\text{var}} - \langle N_{\text{part}} \rangle$ linear dependence
- system size independent
 - the slopes at RHIC and LHC the same
 - a small difference in the offset of ~ 10 fm²



- Peripheral collisions - large Corona percentage
- system size independent
- Central collisions - low Corona percentage
- large system size dependence

Why $5 \text{ GeV}/c < p_T < 8 \text{ GeV}/c$ range?

$$R_{AA}^N = \frac{[\frac{d^2N}{d\eta dp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]_{cen}}{[\frac{d^2N}{d\eta dp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]_{pp,MB}}$$

$$R_{CP}^N = \frac{[\frac{d^2N}{d\eta dp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]_{cen}}{[\frac{d^2N}{d\eta dp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]_{peripheral}}$$

charged particles





- maximum suppression for $5 \text{ GeV}/c < p_T < 8 \text{ GeV}/c$
- the region in p_T for maximum suppression remains the same at:
 - different collision energies
 - different centralities

R_{AA} and R_{AA}^{core} - $\langle N_{part} \rangle$ scaling

$$R_{AA} = \frac{\left(\frac{d^2N}{d\eta dp_T}\right)^{cen}}{\langle N_{bin} \rangle \cdot \left(\frac{d^2N}{d\eta dp_T}\right)^{pp,MB}}$$

$$R_{AA}^{core} = \frac{\left(\frac{d^2N}{d\eta dp_T}\right)^{cen,core}}{\langle N_{bin}^{core} \rangle \cdot \left(\frac{d^2N}{d\eta dp_T}\right)^{pp,MB}}$$





- PHOBOS Collaboration, PRL 96(2006)212301 $\langle N_{part} \rangle$
- STAR Collaboration, PRL 91(2003)172302
- ALICE Collaboration, Phys.Lett.B, 788(2019)166
- ALICE Collaboration, JHEP, 1811(2018)013
- PHENIX Collaboration, Phys. Rev. C69(2004)034910

- R_{AA} scales with $\langle N_{part} \rangle$ for the top RHIC energy
 - R_{AA} scales at LHC energies, although the difference in collision energies is up to a factor 2





- suppression increases by $\sim 20\%$ at low values of $\langle N_{part} \rangle$
 - the difference relative to R_{AA} is decreasing with $\langle N_{part} \rangle$
 - R_{AA}^{core} for very central Cu-Cu and Xe-Xe has the same value as R_{AA}^{core} for Au-Au and Pb-Pb, respectively, at the same $\langle N_{part} \rangle$

R_{AA} and R_{AA}^{core} - $\langle dN_{ch}/d\eta \rangle$ scaling



R_{AA} and R_{AA}^{core} scale with $\langle dN_{ch}/d\eta \rangle$ for all systems and collision energies


$\langle N_{part} \rangle$ versus $\langle dN_{ch}/d\eta \rangle$ scaling



- S_{\perp}^{var} - the same for different systems
- the transverse shapes are similar
- the main difference is in $\langle dN/dy \rangle / S_{\perp}$

- Different shape, size and $\langle dN/dy \rangle / S_{\perp}$ for a given $\langle dN_{ch}/d\eta \rangle$ \Rightarrow their relative contribution to suppression is difficult to unravel

$(1-R_{AA})$ and $\langle dN/dy \rangle/S_{\perp}$ as a function of $\langle N_{part} \rangle$



- For $\langle N_{part} \rangle = 200$, the differences in:
 - $\langle dN/dy \rangle/S_{\perp}$ for Pb-Pb at $\sqrt{s_{NN}} = 2.76, 5.02$ TeV and for Xe-Xe at 5.44 TeV relative to Au-Au at 200 GeV are:
 - 5.25 ± 1 ; 6.77 ± 1 ; 7.89 ± 1 part/fm²
 - $(1-R_{AA})$ values are:
 - 0.10 ± 0.03 ; 0.11 ± 0.03 ; 0.11 ± 0.03 .
- => suppression saturation at LHC energies



- For central Au-Au collisions - $\langle N_{part} \rangle = 350$, the difference in $\langle dN/dy \rangle/S_{\perp}$ between Pb-Pb at 2.76 TeV and Au-Au at 200 GeV is 7 ± 1 part/fm² while the difference in $(1-R_{AA})$ is 0.08 ± 0.03 .

- If $R_{AA} \sim 1 - k \cdot x^2 \cdot T^3$
 - $x^2 \sim S_{\perp}$
 - $T^3 \sim \langle dN/dy \rangle/S_{\perp}$
- => $k^{(2.76 \text{ TeV})} \simeq (0.48 \pm 0.03) \cdot k^{(200 \text{ GeV})}$

- M. Djordjevic et al., Phys. Rev. C99(2019)061902
- B. Betz and Miklos Gyulassy, JHEP 08(2014)090

$R_{AA}^N - \langle N_{part} \rangle$ and $\langle dN_{ch}/d\eta \rangle$ scaling

$$R_{AA}^N = \frac{[\frac{d^2N}{d\eta dp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]^{cen}}{[\frac{d^2N}{d\eta dp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]^{pp,MB}}$$



Suppression in p-p collisions - high multiplicity (HM):

- $\langle \beta_T \rangle$ vs. $[\langle dN/dy \rangle / S_{\perp}]^{1/2}$ scaling

for $[\langle dN/dy \rangle / S_{\perp}]^{1/2} = 3.3 \pm 0.1$



- $S_{\perp}^{pp(HM)} = 7.43 \pm 0.48 \text{ fm}^2$ and $S_{\perp}^{PbPb} = 70 \pm 0.4 \text{ fm}^2$

- assuming the same jet-medium coupling

M. Petrovici et al. Phys.Rev.C, 98(2018)024904

$$\Rightarrow (1 - R_{pp}^{N(HM)}) / (1 - R_{AA}^{N(\langle N_{part} \rangle = 125)}) \approx 0.01 \pm 0.01$$

$(1-R_{AA})/\langle dN/dy \rangle$ and $(1-R_{AA}^N)/\langle dN/dy \rangle - (\langle dN/dy \rangle / S_{\perp})^{1/3}$ dependence



$$\frac{1 - R_{AA}}{\langle dN/dy \rangle} = e^{\alpha - \beta \cdot (\langle dN/dy \rangle / S_{\perp})^{1/3}}$$

- The exponential decrease is similar with the k(T) dependence used in order to reproduce the nuclear modification factors at RHIC and LHC energies (see reference below)



- For $(\langle dN/dy \rangle / S_{\perp})^{1/3} > 2.1$ - constant value, similar with the impact parameter independence of the jet quenching parameter (see reference below)

- C.Andres et al. Nucl. and Part. Phys. Proceedings, 00:1(2018)
 - M.Xie et al. Eur.Phys.J., C79(2019)589

$R_{CP}^N - \langle N_{part} \rangle$ dependence

$$R_{CP} = \frac{[\frac{d^2N}{d\eta dp_T} / \langle N_{bin} \rangle]^{cen}}{[\frac{d^2N}{d\eta dp_T} / \langle N_{bin} \rangle]^{peripheral}}$$

$$R_{CP}^N = \frac{[\frac{d^2N}{d\eta dp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]^{cen}}{[\frac{d^2N}{d\eta dp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]^{peripheral}}$$





- R_{CP}^N scales with $\langle N_{part} \rangle$ for all heavy systems and all collision energies
- the linear dependence as a function of $\langle N_{part} \rangle$ follows from the linear dependence of R_{AA}^N

Collision energy dependence

charged particles

$$R_{CP} = \frac{[\frac{d^2N}{d\eta dp_T} / \langle N_{bin} \rangle]^{cen}}{[\frac{d^2N}{d\eta dp_T} / \langle N_{bin} \rangle]^{peripheral}}$$


$$R_{CP}^N = \frac{[\frac{d^2N}{d\eta dp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]^{cen}}{[\frac{d^2N}{d\eta dp_T} / \langle \frac{dN_{ch}}{d\eta} \rangle]^{peripheral}}$$




STAR Collaboration,
Quark Matter 2012 Conference Proceedings

R_{CP} and R_{CP}^N ($4\text{ GeV}/c < p_T < 6\text{ GeV}/c$) - $\sqrt{s_{NN}}$ dependence (0-5%)/(60-80%)


charged particles



charged particles





π^0



$$R_{CP} \propto a + \frac{b}{s_{NN}} + c \cdot \sqrt{s_{NN}}$$

- PHENIX Collaboration, PRL 109(2012)152301
- PHENIX Collaboration, PRL 101(2008)232301
- ALICE Collaboration, Eur.Phys.J., C74(2014)3108
- ALICE Collaboration, POS(Hard Probes)073

$R_{AA}^{\pi^0}$ and $(1 - R_{AA}^{\pi^0}) / (\langle dN/dy \rangle / S_{\perp}) - \sqrt{s_{NN}}$ dependence



- a maximum in $(1 - R_{AA}^{\pi^0}) / (\langle dN/dy \rangle / S_{\perp})$ at RHIC energies is evidenced followed by a decrease towards LHC energies

- is this a signature of a new state of deconfined matter produced at LHC energies?

- magnetic plasma of light monopoles near T_c




- quarks and gluons dominated deconfined matter

Outlook

- Charged particles R_{AA} , R_{AA}^N , R_{CP} and R_{CP}^N for Au-Au (Cu-Cu) at the top RHIC energy and Pb-Pb (Xe-Xe) at LHC energies, for $5 \text{ GeV}/c < p_T < 8 \text{ GeV}/c$, as a function of $\langle N_{part} \rangle$ and $\langle dN_{ch}/d\eta \rangle$ were discussed
- Considerations based on $1-R_{AA}$ and $\langle dN/dy \rangle/S_{\perp}$ dependence on $\langle N_{part} \rangle$:
 - suppression saturation at LHC energies
 - $k^{LHC} \simeq (0.48 \pm 0.03) \cdot k^{RHIC}$
 - $(1 - R_{pp}^{N(HM)}) / (1 - R_{AA}^{N(\langle N_{part} \rangle \geq 125)}) \simeq 0.01 \pm 0.01$
- R_{CP}^N scales with $\langle N_{part} \rangle$ for all heavy systems and all collision energies
- $(1 - R_{AA}) / \langle dN/dy \rangle - (\langle dN/dy \rangle / S_{\perp})^{1/3}$ exponential dependence
 $(1 - R_{AA}^N) / \langle dN/dy \rangle$ independent on $(\langle dN/dy \rangle / S_{\perp})^{1/3}$ for $(\langle dN/dy \rangle / S_{\perp})^{1/3} > 2.1 \text{ part}/\text{fm}^{2/3}$
- Collision energy dependence of R_{CP} and R_{CP}^N ($4 \text{ GeV}/c < p_T < 6 \text{ GeV}/c$)
 - evidence for saturation at LHC energies
- A maximum in $(1 - R_{AA}^{\pi^0}) / (\langle dN/dy \rangle / S_{\perp})$ at RHIC energies is followed by a decrease towards LHC energies
 - signature for a new state of deconfined matter produced at LHC energies?

Back-up slides



A. Bzdak, B. Schenke, P. Tribedy and R. Venugopalan, Phys.Rev. C87(2013)064906

$$S_{\perp}^{pp} = \pi R_{pp}^2 \quad R_{pp} = l_{fm} f_{pp} - \text{maximal radius for which the energy density of the Yang-Mill fields is larger than } \epsilon = \alpha \Lambda_{QCD}^4 \quad (\alpha \in [1, 10])$$

$$\alpha=1 \quad f_{pp} = \begin{cases} 0.387 + 0.0335x + 0.274x^2 - 0.0542x^3 & \text{if } x < 3.4 \\ 1.538 & \text{if } x \geq 3.4 \end{cases}$$


$$x = (dN_g/dy)^{1/3}$$

$$dN_g/dy \approx dN/dy$$

McLarren, M. Praszalowicz and B. Schenke, Nucl.Phys. A916(2013)210


$$\alpha=10 \quad f_{pp} = \begin{cases} -0.018 + 0.3976x + 0.095x^2 - 0.028x^3 & \text{if } x < 3.4 \\ 1.17 & \text{if } x \geq 3.4 \end{cases}$$

$\langle p_T \rangle$ vs. $[(dN/dy)/S_{\text{perp}}]^{1/2}$



ALICE Collaboration, Nucl.Phys. A931(2014)c888

The slope of $\langle p_T \rangle = f(\text{mass})$ vs. $[(dN/dy)/S_{\text{perp}}]^{1/2}$



$$\text{Slope}_{\langle p_T \rangle = f(\text{mass})} = \alpha + \beta \left(\sqrt{\frac{dN}{dy} / S_{\perp}^{\text{geom}}} \right)^{\gamma}$$

$\langle \beta_T \rangle$ from BGBW fits vs. $[(dN/dy)/S_{\text{perp}}]^{1/2}$

